

ALASKA  
**COVID-19**  
**VACCINE TASK FORCE**

## MODERNA – Vaccine Information

<b>Direct ship size</b>	10 vials per box <ul style="list-style-type: none"> <li>10 dose multi-dose vial, one box has 100 doses (potentially 11 doses per vial)</li> </ul>
<b>Direct ship</b>	Moderna vaccine is shipped at frozen temperatures (-25° to -15°C) pack out with frozen gel packs.
<b>Redistribution</b>	<p>Moderna vaccine can be redistributed at 2-8°C for less than <b>3 hours</b> transit via plane and less than 12 hours transit time via car. If combined, there can be more than 12 total hours of transit time (9 hours via car and 3 hours via plane).</p> <p>Moderna vaccine can be redistributed at -15° to -25°C. There is no limit on transit time. The vaccine must be transported in a transport cooler that can maintain -15° to -25°C with a continuous temperature monitoring device.</p>
<b>Frozen storage</b>	Moderna vaccine can be stored at frozen temperatures (-25° to -15°C) for up to 6 months.
<b>Refrigerated storage</b>	Moderna vaccine can be stored at refrigerator temperatures (2° to 8°C) for up to 30 days. <b>DO NOT REFREEZE A THAWED VIAL</b>
<b>Room temperature</b>	Moderna vaccine can be stored at 15° to 25°C (59° to 77°F) for 12 hours.
<b>Thaw time</b>	<p>Thaw in refrigerated conditions between 2-8°C for 2 hours. Let the vial stand at room temperature for 15 min before use.</p> <p>Thaw at room temperature between 20° to 25°C for 1 hour.</p>
<b>Reconstitution</b>	No dilution required.
<b>Handling open vials</b>	Once opened (needle-punctured), store vial at 2° to 25°C (36 to 77°F) for a maximum of 6 hours.
<b>Dose size</b>	0.5 mL
<b>Injection route</b>	IM
<b>Minimum interval</b>	<p>28 days between dose 1 and dose 2</p> <ul style="list-style-type: none"> <li>Administration of 2nd dose within 4-day grace period (day 17-21) considered valid.</li> <li>If &gt;28 days since 1st dose, 2nd dose should be administered at earliest opportunity (no doses need to be repeated).</li> </ul>
<b>Coadministration with other vaccines</b>	<p>Moderna vaccine should be administered alone with a minimum interval of 14 days before or after administration with any other vaccines.</p> <p>If Moderna vaccine is inadvertently administered within 14 days of another vaccine, doses do not need to be repeated for either vaccine.</p>
<b>Additional information</b>	<ul style="list-style-type: none"> <li>Protect vaccine from light.</li> </ul>