
Read the information on the prescription and label and put the right information into the list.

- You won't need all of the information on each label for the medicine list.
- You may not find some information on the prescription or label. Plan to ask the doctor or pharmacist questions to get the answers.



Right Price Pharmacy

345 W. 35th St, Miami Fl. 33165

Phone (305) 249-5598

Dr. Hency, Robert B

Sandra Birdsnest ALBUTEROL .83 MG/ML

Equivalent to PROVENTIL .83 MG/ML 2 PUFFS TWICE A DAY

QTY 3.0 05/09/112 Discard after 05/09/14 0 Refills/Dr. approval reg'd/

NEW RX

Therapeutic substance in drug

When not to use

stop taking it,

when to see a

doctor, and

effects

possible side

this drug, when to

Drug Facts

Active ingredient (in each tablet) Purpose
Chlorpheniramine maleate 2 mg Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allerigies:

- sneezing
- runny nose
- itchy, watery eyes
- itchy throat

Warnings

Product type

Symptoms or diseases the drug treats

Ask a doctor before use if you have ■ glaucoma

- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives

When using this procuct

- You may get drowsy
- Avoid alcoholic drinks
- Alcohol, sedatives, and tranquilizers may increase drowsiness
- Be careful when driving a motor vehicle or operating machinery
- Excitability may occur, especially in children

If pregnant or breastfeeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years and over	Take 2 tablets every 4 to 6 hours; not more than 12 tablets in 24 hours
Children 6 years to under 12 years	Take 1 tablet every 4 to 6 hours; not more than 6 tablets in 24 hours
Children under 6 years	Ask a doctor

Read carefully: how much to take, how often to take it, and when to stop taking it

More information on how to store the drug

Other information Store at 20-25° C (68-77° F)

■ Protect from excessive moisture

Inactive ingredients D&C yellow no. 10, lactose, magnesium stearate, microcrystalline cellulose, pregelatinized starch Other things in the drug, such as colors or flavorings

BigPill

EXTRA STRENGTH

Antacid

Fast-acting Heartburn Relief

Long-lasting Antacid Barrier
Original Flavor
100 CHEWABLE
TABLETS

Drug Facts

Uses relieves

- Acid indigestion * heartburn * sour stomach
- Upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have * kidney disease * a magnesium or sodium restricted diet

Ask a doctor before use if you are now taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product do not take more than 16 tablets in a 24 hour period or use the maximum dosage for more than 2 weeks

Keep out of reach of children

Directions

- Chew 2 to 4 tablets four times a day or as directed by a doctor
- Take after meals and at bedtime or as needed
- For bet results follow by a half glass of water or other liquid
- Do not swallow whole

Other information

- Each tablet contains: magnesium 50 mg and sodium 32 mg
- Store at 20⁰-25⁰ C

Inactive ingredients alginic acid, calcium stearate, flavor, mannitol,

Distributed by Redrobin & Co. Paris, Texas 22904 Health



